



t

Nutritional and Health-Related Environmental Studies Section

Technical Meeting on Environmental Enteric Dysfunction, Microbiome and Undernutrition

IAEA, Vienna, Austria

October 28-30, 2015

Scientific Secretary: Ms. Cornelia Loechl, Nutrition Specialist

Phone: +43 (1) 2600-21635

Email: C.Loechl@iaea.org

Despite progress made to date, a large proportion of the population in low- and middle-income countries (LMICs) still live in environments characterized by poor water, sanitation and hygiene (WASH) conditions. According to the WHO and UNICEF, an estimated 2.5 billion people remain without access to improved sanitation facilities (2015)¹. In addition, many LMICs continue to struggle with a high burden of undernutrition, demonstrated by the nearly 805 million undernourished people worldwide, the vast majority of which live in LMICs (2012-2014)². Environmental enteric dysfunction (EED), a subclinical inflammatory disorder of the gut, is also highly common among impoverished inhabitants of environments with poor sanitation and hygiene, such as those often found in LMICs, and has gained increasing interest as a key pathway linking unsanitary living conditions with different forms of undernutrition (stunting, wasting, underweight, etc.)³⁻¹⁰. It is hypothesized that the failure of health interventions, specifically targeting malnutrition in LMICs, can be directly attributed to EED¹¹. Likewise, oral vaccines have been observed as being less immunogenic in children from LMICs suggesting that EED may cause altered mucosal immunity in these children¹¹.

EED is the result of recurring infection in the small intestine via the faecal-oral ingestion route. Constant exposure to commensal and/or pathogenic bacteria in the small intestine induce inflammatory processes, partial villous atrophy, and crypt hyperplasia through a T-cell mediated process, resulting in increased intestinal permeability, decreased nutrient absorption, bacterial translocation, and innate and acquired immune system activation¹². However, individuals affected typically remain asymptomatic – rarely having overt gastrointestinal symptoms or diarrhoea¹¹. These EED-related histological changes of the gut, though, result in harmful health effects including malabsorption and ineffective utilisation of nutrients, as well as a blunted innate immune response, potentially leading to oral vaccine failure and increased susceptibility to infection¹¹. One of the most significant impacts of EED may be on undernutrition and potentially growth faltering in children⁵. Moreover, EED potentially contributes to poor response to nutrition therapy¹¹. This may ultimately increase child morbidity and mortality, reduce cognitive development, and reduce adult economic productivity.

Multiple studies are underway to explore the complex relationship between WASH, EED, and child malnutrition, health and development¹³⁻¹⁵. While this condition is still poorly understood, there is a growing body of evidence that suggests the intermediate biological mechanisms of undernutrition and EED are interrelated in a complex manner – acting both synergistically and as part of a feedback system. A chronically damaged and inflamed gut may increase the likelihood of developing a bodily predisposition to becoming undernourished through several pathways; while being undernourished further compromises healthy intestinal lining, thus increasing susceptibility to additional infection. Chronic inflammation, a symptom of EED, has been associated with altered absorption and ineffective utilisation of micronutrients and macronutrients. Likewise, repetitive infection causing chronic immune activation may result in absorbed nutrients (e.g. amino acids) being diverted towards inflammatory processes instead of being used as building blocks or for serving their metabolic functions¹⁶.

Furthermore, malabsorption may lead to perturbed micronutrient homeostasis, as in the case of iron¹⁷, which can result in anaemia and suboptimal organ development in children. Iron deficiency can also disturb intestinal permeability in children¹⁸. Vitamin A and zinc are crucial for maintaining optimal gut health in early life¹⁹, and may be influential to preventing and treating EED. Vitamin A preserves gut integrity and is associated with increased permeability of the mucosal lining and increased susceptibility to enteric infection under deficient conditions²⁰⁻²³. Fluctuations in zinc homeostasis may develop when reduced absorptive capacity, resulting from damaged intestinal mucosa, occurs due to EED^{24, 25}. This zinc imbalance both further exacerbates intestinal abnormalities and perpetuates EED through several pathways, including higher susceptibility to enteric infections, disturbed barrier function, and the onset of chronic systemic inflammation²⁶.

The gut microbiome is subject to a growing trend in scientific interest – it plays vital physiological, immunological, and nutritional/metabolic roles in humans, and has been recognised as one of the primary mediators in several pathological conditions including EED²⁷. Small intestine bacterial

overgrowth (SIBO), a disorder marked by symptoms including diarrhoea, malnutrition, weight loss and malabsorption, is associated with nutritional deficiencies and should be considered as an additional component to EED linking unsanitary living conditions to child undernutrition and growth faltering^{28,29}. Numerous studies have revealed that specific relationships exist between the gut microbiome and both human metabolism and body composition in healthy individuals, but their relevance to individuals with EED warrants further investigation³⁰. Indeed, children diagnosed with inflammatory intestinal disorders – similar to EED - have altered body composition (the relative amounts of fat and lean tissue) and show a decrease in lean tissue, which can persist over time³¹. Also, in stunted children, often suffering from EED, a greater accumulation of body fat and a lower gain in lean mass was observed³².

A clear gold standard for diagnosis of EED is not yet defined. The most widely used surrogate marker for epithelial integrity is the lactulose: mannitol test, in which the subject is given an oral solution containing lactulose and mannitol. Several serum and faecal biomarkers have been studied as indicators of intestinal inflammation with a few having only been studied in populations at risk for EED¹¹. In this regard, more research is needed to assess potential biomarkers for the diagnosis of EED.

Considering the bidirectional relationship between EED and undernutrition, holistic and targeted nutrition interventions (prevention and treatment) are needed to concurrently mitigate the histological and functional changes associated with EED while correcting for nutritional deficiencies³³. Improving WASH conditions and knowledge, attitudes, and practices (KAPs) is undoubtedly of great importance when trying to restore intestinal functionality. However, solely improving WASH conditions and KAPs is not sufficient to combat EED or mitigate subsequent child growth faltering, as demonstrated in Nepal³⁴. Vitamin and mineral supplementation should be considered as a viable intervention as multiple studies have shown that iron, vitamin A, and zinc supplementation – given individually (vitamin A, zinc) or in combination (iron, vitamin and zinc) – ameliorate intestinal morphology, decrease intestinal permeability, improve gut functionality, and promote linear growth^{20,21, 27, 35, 36}. However, nutritional status and intakes of specific micronutrients may also influence the gut microbiome. It appears that an increase in unabsorbed dietary iron through fortification or supplementation could modify the colonic microbiota equilibrium and favour growth of pathogenic strains over beneficial ‘barrier’ strains. Additionally, studies have revealed that providing micronutrient powders with iron to weaning infants adversely affects the gut microbiome, increasing pathogen abundance and causing intestinal inflammation⁴²⁻⁴⁸.

The goal of the International Atomic Energy Agency (IAEA) is to enhance the capabilities of Member States to address needs related to the management of all forms of malnutrition. The IAEA contributes towards the global health nutrition agenda by promoting the use of stable isotope techniques in developing and evaluating nutrition interventions, which are more accurate and sensitive than conventional techniques, safe to use across all ages, and can be used in a community setting. Supporting the use of stable isotope techniques to assess the bioavailability of micronutrients – including iron and zinc – from foods and assessing an individuals’ vitamin A status could provide Member States with the necessary evidence to design or improve the management of EED and undernutrition, especially as micronutrients have been demonstrated as having a function in both exacerbating (deficiency) and treating (adequate intake) EED. Stable isotope techniques have, in fact, already been identified by an expert group as the method of choice for evaluating the role of zinc nutrition in EED vulnerable children³⁷. Furthermore, these techniques were recognized as a potent tool in investigating micronutrient absorption in children with EED, and as such, they were included in the report of the Second World Congress of Pediatric Gastroenterology, Hepatology and Nutrition⁴¹. Stable isotopes can also be used to assess body composition (deuterium dilution technique), which could be crucial in investigating the impact EED has on the quality of growth in early life. There is also emerging evidence that demonstrates the potential use of breath tests with ¹³C-labelled sugars (sucrose and glucose) as a novel, non-invasive biomarker of gut integrity and absorptive capacity to diagnose children with EED and detect SIBO^{38,39}. Stable isotopes can also be

effectively used to measure the metabolic activity of the gut microbiome, which could contribute towards revealing the microbiome's influence on EED and undernutrition⁴⁰.

Keeping in mind the promising role of stable isotope techniques to contribute to a better understanding of the relationships between EED, the microbiome and undernutrition, the IAEA plans to organize a Technical Meeting to discuss the multifaceted and interrelated challenges and research gaps in the area of EED.

Meeting Objectives

This technical meeting will take place over the course of three days, 28-30 October 2015, and bring together experts working in the field of EED and experts knowledgeable in the application of stable isotope techniques in nutrition. The meeting will have the following objectives:

- Discuss current knowledge and gaps on the causes and consequences of EED
- Share experiences related to the implementation and evaluation of programmes to prevent and treat EED in infants, children, and adults
- Discuss technical issues and strategies related to the management of EED and undernutrition, including tests for diagnosis
- Identify knowledge gaps in the field of EED where the IAEA can add value by supporting the use of stable isotope techniques

List of Topics

- Interactions between EED and undernutrition
- Nutrition interventions in children and adults suffering from EED and undernutrition
- Effects of EED on growth, body composition and functional outcomes
- Application of novel, low-cost, non-invasive methods of diagnosis of inflammation, permeability and SIBO with a focus on stable isotope techniques
- Contribution of stable isotope techniques to assess absorption of micronutrients (zinc, iron, and vitamin A) in the context of EED

Expected outcome

The outcome of the Technical Meeting will be recommendations that identify the role of the IAEA in the global effort to address EED and undernutrition through the application of stable isotope techniques.

References

1. UNICEF. Water, Sanitation, and Hygiene (WASH) Press centre. (http://www.unicef.org/media/media_45481.html)
2. FAO, IFAD, WFP. The State of Food Insecurity in the World 2014. (<http://www.fao.org/publications/sofi/en/>)
3. Humphrey JH. (2009). Child undernutrition, tropical enteropathy, toilets, and handwashing. *Lancet* 374:1032-35.
4. Lin A, Arnold BF, Afreen S, Goto R, Nurul Huda TM, Haque R, Raqib R, Unicomb L, Ahmed T, Colford Jr JM, Luby SP. (2013). Household environmental conditions are associated with enteropathy and impaired growth in rural Bangladesh. *Am J Trop Med Hyg* 89(1):1230-7.
5. Campbell DI, Lunn PG, Elia M. (2002). Age-related association of small intestinal mucosal enteropathy with nutritional status in rural Gambian children. *Br J Nutr* 88:499-505.
6. Lunn PG, Northrop-Clewes CA, Downes RM. (1991). Intestinal permeability, mucosal injury, and growth faltering in Gambian infants. *Lancet* 338:907-910.
7. Keusch GT, Rosenberg IH, Denno DM, Duggan C, Guerrant RL, Lavery JV, Tarr PI, Ward HD, Black RE, Nataro JP, Ryan ET, Bhutta ZA, Coovadia H, Lima A, Ramakrishna B, Zaidi AKM, Hay Burgess DC, Brewer T. (2013). Implications of acquired environmental enteric dysfunction for growth and stunting in infants and children living in low- and middle-income countries. *FNB* 34(3):357-63.
8. Hossain MI, Nahar B, Hamadani JD, Ahmed T, Roy AK, Brown KH. (2010). Intestinal mucosal permeability of severely underweight and nonmalnourished Bangladeshi children and effects of nutritional rehabilitation. *J Pediatr Gastroenterol Nutr* 51:638-44.
9. Brewster DR, Manary MJ, Menzies IS, O'Loughlin EV, Henry RL. (1997). Intestinal permeability in Kwashiorkor. *Arch Dis Child* 76(3):236-41.
10. Jones KDJ, Huenten-Kirsch B, Laving AMR, Munyi CW, Ngari M, Mikusa J, Mulongo MM, Odera D, Nassir HS, Timbwa M, Owino M, Fegan G, Murch SH, Sullivan PB, Warner JO, Berkley JA. (2014). Mesalazine in the initial management of severely acutely malnourished children with environmental enteric dysfunction: a pilot randomized controlled trial. *BMC Medicine* 12:133.
11. Korpe PS, Petri WA. (2012). Environmental Enteropathy: Critical implications of a poorly understood condition. *Trends Mol Med* 18(6):328-36.
12. Ramakrishna BS, Venkataraman S, Mukhopadhyaya A. (2006). Tropical malabsorption. *Postgrad Med J* 82:779-87.
13. MAL-ED. The Interactions of Malnutrition & Enteric Infections: Consequences for Child Health and Development. (<http://mal-ed.fnih.org/>)
14. WASH Benefits Study: Bangladesh and Kenya. (<http://www.washbenefits.net/>)
15. SHINE: Sanitation, Hygiene, Infant Nutrition Efficacy Project. (<http://clinicaltrials.gov/show/NCT01824940>)
16. Gautron L, Sophie L. (2009). Neurobiology of inflammation-associated anorexia. *Front Neurosci* 3:59.
17. Weiss G, Goodnough LT. (2005). Anemia of chronic disease. *N Engl J Med* 352:1011-1023.
18. Berant M, Khourie M, Menzies IS. (1992). Effect of iron deficiency on small intestinal permeability in infants and young children. *J Pediatr Gastroenterol Nutr* 14(1):17-20.
19. Ziegler TR, Evans ME, Fernandez-Estivariz C, Jones DP. (2003). Trophic and cytoprotective nutrition for intestinal adaptation, mucosal repair, and barrier function. *Annu Rev Nutr* 23:229-61.
20. Thurnham DI, Northrop-Clewes CA, McCollough FS, Das BS, Lunn PG. (2000). Innate immunity, gut integrity, and vitamin A in Gambian and Indian infants. *J Infect Dis* 182(1):23-8.
21. Chen P, Soares M, Lima AA, Gamble MV, Schorling JB, Conway M, Barrett LJ, Blaner WS, Guerrant RL. (2003). Association of vitamin A and zinc status with altered intestinal permeability: analyses of cohort data from northeastern Brazil. *J Health Popul Nutr* 21(4):309-15.
22. Curtale F, Pokhrel RP, Tilden RL, Higashi G. (1995). Intestinal helminths and xerophthalmia in Nepal. A case control study. *J Trop Pediatr* 41:334-337.
23. Sommer A, Tarwotjo I, Katz J. (1987). Increased risk of xerophthalmia following diarrhea and respiratory disease. *Am J Clin Nutr* 45:977-980.
24. Manary MJ, Hotz C, Krebs NF, Gibson RS, Westcott JE, Broadhead RL, Hambidge KM. (2002). Zinc homeostasis in Malawian children consuming a high-phytate, maize-based diet. *Am J Clin Nutr* 75:1057-61.
25. Manary MJ, Abrams MS, Griffin IJ, Quimper MM, Shulman RJ, Hamzo MG, Chen Z, Maleta K, Manary MJ. (2010). Perturbed zinc homeostasis in rural 3-5-y-old Malawian children is associated with abnormalities in intestinal permeability attributed to tropical enteropathy. *Pediatr Res* 67:671-5.
26. Lindenmayer GW, Stoltzfus RJ, Prendergast AJ. (2014). Interactions between zinc deficiency and environmental enteropathy in developing countries. *Adv Nutr* 5:1-6.

27. Kau AL, Ahern PP, Griffin NW, Goodman AL, Gordon JI. (2012). Human nutrition, the gut microbiome, and immune system: envisioning the future. *Nature* 474(75351): 327-336.
28. Fagundes Neto U, Martins MC, Lima FL, Patricio FR, Toledo MR. (1994). Asymptomatic environmental enteropathy among slum-dwelling infants. *J Am Coll Nutr* 13(1): 51-6.
29. Donowitz JR, Petri WA Jr. (2014). Pediatric small intestinal bacterial overgrowth in low-income countries. *Trends in Mol Med*: doi:10.1016/j.molmed.2014.11.001
30. Vrieze A, Holleman F, Zoetendal EG, de Vos WM, Hoekstra JBL, Nieuwdorp M. (2010). The environment within: how gut microbiota may influence metabolism and body composition. *Diabetologia* 53: 606-613.
31. Hill RJ. (2014). Update on nutritional status, body composition and growth in paediatric inflammatory bowel disease. *World J Gastroenterol* 20(12): 3191-3197.
32. Martins PA, Hoffman DJ, Fernandes MT, Nascimento CR, Roberts CR, Sesso R, Sawaya AL. (2004). Stunted children gain less lean body mass and more fat mass than their non-stunted counterparts: a prospective study. *Br J Nutr* 92(5):819-25.
33. McKay S, Gaudier E, Campbell DI, Prentice AM, Albers R. (2010). Environmental enteropathy: new targets for nutritional interventions. *Int Health* 2:172-180.
34. Langford R, Lunn P, Panter-Brick C. (2011). Hand-washing, subclinical infections, and growth: a longitudinal evaluation of an intervention in Nepali slums. *Am J Hum Biol* 23:621-629.
35. Ryan KN, Stephenson KB, Trehan I, Shulman RJ, Thakwalakwa C, Murray E, Maleta K, Manary MJ. (2014). Zinc or albendazole attenuates the progression of environmental enteropathy: a randomized controlled trial. *Clin Gastroenterol Hepatol* 12(9):1507-13.
36. Smith HE, Ryan KN, Stephenson KB, Westcott C, Thakwalakwa C, Maleta K, Cheng JY, Brenna JT, Shulman RJ, Trehan I, Manary MJ. (2014). Multiple micronutrient supplementation transiently ameliorates environmental enteropathy in Malawian children aged 12-35 months in a randomized controlled trial. *J of Nutr* 144(12):2059-65.
37. Young GP, Mortimer EK, Gopalsamy GL, Alpers DH, Binder HJ, Manary MJ, Ramakrishna BS, Brown IL, Brewer TG. (2014). Zinc deficiency in children with environmental enteropathy-development of new strategies: report from an expert workshop. *Am J Clin Nutr* 100:1198-207.
38. Ritchie BK, Brewster DR, Davidson GP, Tran CD, McNeil Y, Hawkes JS, Butler RN. (2009). ¹³C-Sucrose breath test: Novel use of a non-invasive biomarker of environmental gut health. *Pediatrics* 124(2):620-626.
39. Banik GD, Maity A, Som S, Ghosh C, Daschakraborty SB, Chaudhuri S, Ghosh S, Pradhan M. (2014). Diagnosis of small intestinal bacterial overgrowth in irritable bowel syndrome patients using high-precision stable ¹³CO₂/¹²CO₂ isotope ratios in exhaled breath. *J Anat At Spectrom* 29:1918-1924.
40. Reichardt N, Barclay AR, Weaver LT, Morrison DJ. (2011). Use of stable isotopes to measure the metabolic activity of the human intestinal microbiota. *Appl Environ Microbiol* 77(22):8009-8014.
41. Allen S, Brewster DR, Elliot EJ, Fasano A, Phillips AD, Sanderson IR, Phillip I. (2004). Intestinal infections and environmental enteropathy: working group report of the Second World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr* 39(2):662-9.
42. Andrews SC, Robinson AK, Rodriguez-Quinones F. (2003). Bacterial iron homeostasis. *FEMS Microbiol Rev* 27:215-37.
43. Naikare H, Palyada K, Panciera R, et al. (2006). Major role for FeoB in *Campylobacter jejuni* ferrous iron acquisition, gut colonization, and intracellular survival. *Infect Immun* 74:5433-44.
44. Anderson RC, Cookson AL, McNabb WC, et al. (2010). *Lactobacillus plantarum* DSM 2648 is a potential probiotic that enhances intestinal barrier function. *FEMS Microbiol Lett* 309:184-92.
45. Yatsunencko. (2012). Human gut microbiome viewed across age and geography. *Nature*.
46. Zimmermann MB, Chassard C, Rohner F, et al. (2010). The effects of iron fortification on the gut microbiota in African children: a randomized controlled trial in Cote d'Ivoire. *Am J Clin Nutr* 92:1406-15.
47. Soofi S, Cousens S, Iqbal SP, et al. (2013). Effect of provision of daily zinc and iron with several micronutrients on growth and morbidity among young children in Pakistan: a cluster-randomised trial. *Lancet* 382:29-40.
48. Jaeggi T, Kortman GA, Moretti D, Chassard C, Holding P, Dostal A, Boekhorst J, Timmerman HM, Swinkels DW, Tjalsma H, Njenga J, Mwangi A, Kvalsvig J, Lacroix C, Zimmermann MB. (2014). Iron fortification adversely affects the gut microbiome, increases pathogen abundance and induces intestinal inflammation in Kenyan infants. *Gut*. Aug 20.